

For Research Use Only

TrkA Monoclonal antibody

Catalog Number: 60548-1-Ig



Basic Information

Catalog Number:

60548-1-Ig

Concentration:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG27985

GenBank Accession Number:

NM_001007792

GeneID (NCBI):

4914

UNIPROT ID:

P04629

Full Name:

neurotrophic tyrosine kinase,
receptor, type 1

Calculated MW:

88 kDa

Observed MW:

140 kDa

Purification Method:

Protein G purification

CloneNo.:

3B11C7

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : Karpas-299 cells, NK-92 cells, SH-SY5Y cells, IMR-32 cells, TF-1 cells, K-562 cells, MOLT-4 cells, SK-N-SH cells

Background Information

NTRK1 (Neurotrophic tyrosine kinase receptor 1) is located on chromosome 1q23.1 and is responsible for a rare genetic disorder, CIPA (congenital insensitivity to pain with anhidrosis) (PMID: 37907480). As a neurotrophin receptor, NTRK1 is abundantly expressed in human neuronal tissues and plays an integral role in the development and function of the human nervous system (PMID: 27843590). NTRK1 also regulates the survival and proliferation of cholinergic neurons (PMID: 20054812).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



NK-92 cells were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 60548-1-Ig (NTRK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.