

For Research Use Only

CD268/BAFF-R Monoclonal antibody

Catalog Number:60763-7-Ig



Basic Information

Catalog Number: 60763-7-Ig	GenBank Accession Number: BC105123	Purification Method: Protein A purification
Source: Mouse	GeneID (NCBI): 115650	CloneNo.: 2B12D1
Isotype: IgG2a	UNIPROT ID: Q96RJ3	Recommended Dilutions: IHC: 1:1000-1:4000
Immunogen Catalog Number: AG18512	Full Name: tumor necrosis factor receptor superfamily, member 13C	
	Calculated MW: 184 aa, 19 kDa	

Applications

Tested Applications: IHC, ELISA	Positive Controls: IHC : human tonsillitis tissue, human spleen tissue
Species Specificity: human	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

B-cell-activating factor receptor (BAFF-R, also known as CD268 and TNFRSF13C) is a member of the tumor necrosis factor receptor (TNFR) superfamily. It is expressed in most malignant B cells, including those in primary central nervous system lymphoma (PCNSL), and is involved in the survival and proliferation of these cells (PMID: 32528875). The interaction of BAFF with BAFF-R promotes NF- κ B activation, which plays a crucial role in B-cell maturation, survival, and costimulation of T-cell activation and proliferation (PMID: 30349534). Additionally, BAFF-R is associated with chronic kidney disease (CKD) by activating NF- κ B signaling in renal tubular epithelial cells (PMID: 38791453).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 60763-7-Ig (CD268/BAFF-R antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).