

For Research Use Only

FMO1 Monoclonal antibody, PBS Only

Catalog Number:61032-3-PBS



Basic Information

Catalog Number: 61032-3-PBS	GenBank Accession Number: NM_001282692	Purification Method: Protein G purification
Source: Mouse	GeneID (NCBI): 2326	CloneNo.: 1B9G6
Isotype: IgG1	UNIPROT ID: Q01740	
Immunogen Catalog Number: AG34286	Full Name: flavin containing monooxygenase 1	
	Calculated MW: 60 kDa	
	Observed MW: 50-53 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse, rat, pig, rabbit

Background Information

Flavin-containing monooxygenase 1 (FMO1) belongs to the FMO family. FMO1 is an enzyme that catalyzes the oxygenation of a series of sulfur-containing and nitrogen-containing substrates, and thus is involved in a series of drugs and xenobiotics metabolism. FMO1 can regulate PPAR- α and mediate metabolic processes. FMO1 is expressed at relatively high levels in human fetal liver and also represents the major FMO enzyme in the adult human and rabbit intestine and kidney (PMID: 11809920, 30757917, 37553313). Human FMO1 has 2 isoforms with the molecular weights of 53 kDa and 60 kDa.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 61032-3-Ig (FMO1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 61032-3-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 61032-3-Ig (FMO1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 61032-3-PBS in a different storage buffer formulation.



pig kidney tissue were subjected to SDS PAGE followed by western blot with 61032-3-Ig (FMO1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 61032-3-PBS in a different storage buffer formulation.