For Research Use Only

Anti-Mouse CD8a (53-6.7) Catalog Number:65069-1-lg 5 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 65069-1-lg	· · · · · · · · · · · · · · · · · · ·		Purification Method: Affinity purification
	Concentration:	GenelD (NC	BI).	CloneNo.:
	500ug, 0.5 mg/ml	12525		53-6.7
	Source: Rat	UNIPROT ID P01731	:	
	Isotype: IgG2a, kappa	Full Name: CD8 antiger	n, alpha chain	
Applications	Tested Applications: FC			
	Cited Applications: FC			
	Species Specificity: mouse			
	Cited Species: mouse			
Background Information	CD8 is a transmembrane glycoprotein composed of two disulfide-linked chains. It can be present as a homodimer of CD8 α or as a heterodimer of CD8 α and CD8 β (PMID: 3264320; 8253791). CD8 is found on most thymocytes. The majority of class I-restricted T cells express mostly the CD8 α β heterodimer while CD8 α α homodimers alone have been found on some gut intraepithelial T cells , on some T cell receptor (TCR) γ δ T cells and on NK cells (PMID: 2111591; 1831127; 8420975). CD8 acts as a co-receptor that binds to MHC class-I and participates in cytotoxic T cell activation (PMID: 8499079). During T cell development, CD8 is required for positive selection of CD4-/CD8+T cells (PMID: 1968084).			
	have been found on some g (PMID: 2111591; 1831127; 8 T cell activation (PMID: 849	gut intraepithelial T cel 3420975). CD8 acts as a	ls , on some T cell rece co-receptor that bind	eptor (TCR) $\gamma^{-\delta}$ T cells and on NK cells s to MHC class-I and participates in cytotoxi
	have been found on some g (PMID: 2111591; 1831127; 8 T cell activation (PMID: 849	gut intraepithelial T cel 3420975). CD8 acts as a	ls , on some T cell rece co-receptor that bind	eptor (TCR) $\gamma \delta$ T cells and on NK cells s to MHC class-I and participates in cytotoxi juired for positive selection of CD4-/CD8+ 1
	have been found on some g (PMID: 2111591; 1831127; 8 T cell activation (PMID: 849 cells (PMID: 1968084).	gut intraepithelial T cel 3420975). CD8 acts as a 99079). During T cell de	s , on some T cell reco co-receptor that bind velopment, CD8 is rec	eptor (TCR) $\gamma^{-\delta}$ T cells and on NK cells s to MHC class-I and participates in cytotoxi
Notable Publications	have been found on some g (PMID: 2111591; 1831127; 8 T cell activation (PMID: 849 cells (PMID: 1968084).	gut intraepithelial T cel 3420975). CD8 acts as a 19079). During T cell de Pubmed ID	s , on some T cell rece co-receptor that bind velopment, CD8 is rec Journal	eptor (TCR) γ δ T cells and on NK cells s to MHC class-I and participates in cytotoxi quired for positive selection of CD4-/CD8+1 Application
	have been found on some g (PMID: 2111591; 1831127; 8 T cell activation (PMID: 849 cells (PMID: 1968084). Author Huajing Wang	gut intraepithelial T cel 3420975). CD8 acts as a 19079). During T cell de Pubmed ID 34090420	s , on some T cell reco co-receptor that bind velopment, CD8 is rec Journal Malar J	eptor (TCR) γ δ T cells and on NK cells s to MHC class-I and participates in cytotox juired for positive selection of CD4-/CD8+ Application FC FC

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^{^6}C 57BL/6 mouse splenocytes were surface stained with CoraLite® Plus 647 Anti-Mouse CD3 (17A2) and 0.5 ug Anti-Mouse CD8a (65069-1-lg, Clone:53-6.7) or 0.5 ug Rat IgG2a Isotype Control (2A3) and FITC anti-rat IgG2a Antibody at dilution 1:100. Cells were not fixed.