For Research Use Only

Anti-Human CD80 (B7-1) (2D10.4)

Catalog Number:65083-1-Ig



Purification Method:

Affinity purification

CloneNo.:

2D10.4

Basic Information

Catalog Number:

65083-1-lg

Size:

100ug, 500 μg/ml Source:

Mouse Isotype: IgG1, kappa GenBank Accession Number:

BC042665 GeneID (NCBI):

Genera (NCB)

33 kDa

Full Name: CD80 molecule Calculated MW:

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide, pH 7.3.

Selected Validation Data



1X10^6 Daudi cells were surface stained with 0.25 ug Anti-Human CD80 (B7-1) (65083-1-1g, Clone: 2D10.4) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), or stained with 0.25 ug isotype control and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (black). Cells were not fixed.