For Research Use Only

Anti-Human CD40 (G28.5)

Catalog Number: 65103-1-Ig



Purification Method:

Affinity purification

CloneNo.:

G28.5

Basic Information

Catalog Number:

65103-1-lg

Size:

100ug, 0.5 mg/ml

Source: Mouse Isotype:

IgG1, kappa

Full Name: CD40 molecule, TNF receptor superfamily member 5

BC012419

958

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG0000101017

GenBank Accession Number:

Calculated MW: 277 aa, 31 kDa

Species Specificity:

Human

Tested Applications:

FC

Background Information

CD40, also named as TNFRSF5 and Bp50, is a type I transmembrane protein that belongs to the tumor necrosis factor-R (TNF-R) family (PMID: 10647992). CD40 is expressed on B cells, dendritic cells (DCs), monocytes, platelets, and macrophages as well as by non-hematopoietic cells such as myofibroblasts, fibroblasts, epithelial, and endothelial cells (PMID: 19426221). It is a receptor for TNFSF5/CD40LG. CD40L/CD40 interactions are essential for immunoglobulin (Ig) isotype switching, germinal center formation and the development of B cell memory (PMID: 7516405; 11675497).

Storage

Applications


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD40 (65103-1-lg, Clone:G28.5) or Mouse IgG1 Isotype Control and CoraLite®647-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were then stained with CoraLite®488 Anti-Human CD19. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD40 (65103-1-1g, Clone:G28.5) or Mouse IgG1 Isotype Control and CoraLite®647-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.