

For Research Use Only

Anti-Mouse TCR beta (H57-597)

Catalog Number: 65106-1-Ig



Basic Information

Catalog Number: 65106-1-Ig	GenBank Accession Number: GeneID (NCBI): 21577	Purification Method: Affinity purification
Size: 100ug, 0.5 mg/ml	Full Name: T-cell receptor beta chain	CloneNo.: H57-597
Source: Armenian Hamster		
Isotype: IgG		

Applications

Tested Applications:
FC

Species Specificity:
Mouse

Background Information

T-cell receptor (TCR) consists of heterodimeric glycoproteins (TCR alpha/beta or gamma/delta) that demonstrate homology with immunoglobulins (PMID: 2144901). TCR beta is a member of the immunoglobulin superfamily and is expressed on alpha/beta TCR-bearing T cells and thymocytes. The TCR heterodimers are noncovalently associated with CD3. The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta (PMID: 1826255). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Storage:
Store at 2-8°C.

Storage Buffer:
Phosphate based buffer with 0.09% sodium azide, pH 7.2.

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ mouse splenocytes were surface stained with 0.5 ug Anti-Mouse TCR beta (65106-1-Ig, Clone:H57-597) and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:200 (green), or stained with 0.5 ug isotype control antibody and FITC Goat anti-hamster (Armenian) IgG Antibody at dilution 1:200 (black). Cells were not fixed.