

For Research Use Only

Anti-Human CD154 (24-31)

Catalog Number: 65156-1-Ig



Basic Information

Catalog Number:

65156-1-Ig

Size:

100ug, 1 mg/ml

Source:

Mouse

Isotype:

IgG1, kappa

GenBank Accession Number:

BC071754

GeneID (NCBI):

959

UNIPROT ID:

P29965

Full Name:

CD40 ligand

Calculated MW:

261 aa, 29 kDa

Purification Method:

Purified by protein-A affinity chromatography

CloneNo.:

24-31

Applications

Tested Applications:

FC

Species Specificity:

Human, Non-human primates

Background Information

The CD40 ligand (CD40L, TRAP, CD154), a member of the TNF superfamily of ligands, is expressed as either a 33-kd transmembrane homologue or 18-kd soluble form (sCD154). CD40L is primarily expressed on activated CD4+ T cells and on a small proportion of CD8+ T cells and platelets. It binds to CD40 on antigen-presenting cells (APC), which leads to many effects depending on the target cell type. It has been suggested that CD40/CD40L interactions regulate oxidative stress and affect various signaling pathways in both the immunological and the cardiovascular systems. The CD40/CD40L system is also involved in tumorigenesis. Its expression is tightly regulated, and abnormal levels of CD40L are associated with the pathogenesis of atheromatous plaque destabilization and thrombotic events. Multiple mutations in CD40LG gene have been identified that are associated with hyper-IgM immunodeficiency syndrome type 1.

Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide.

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 PMA (50 ng/ml) and ionomycin (1 ug/ml)-stimulated human PBMCs (5 hours) were surface stained with 0.2 ug Anti-Human CD154 (65156-1-Ig, Clone: 24-31) and FITC anti-mouse IgG1 Antibody at dilution 1:200 (green), or stained with 0.2 ug isotype control and FITC anti-mouse IgG1 Antibody at dilution 1:200 (black). Cells were not fixed.