For Research Use Only

## Anti-Rat CD11b (OX42) Catalog Number:65229-1-lg




| Basic Information      | Catalog Number:<br>65229-1-lg   | GenBank Accession Number:<br>NM_012711 | Purification Method:<br>Affinity purification |
|------------------------|---|--|---|
|                        | Size:<br>100ug, 0.5 mg/ml   | GenelD (NCBI):<br>25021                | CloneNo.:<br>OX42                             |
|                        | Source:<br>Mouse  | UNIPROT ID:<br>Q9JI30                  |   |
|                        | <mark>Isotype:</mark><br>IgG2a, kappa   | Full Name:<br>integrin alpha M         |   |
|                        |   | Calculated MW:<br>127 kDa              |   |
| Applications           | Tested Applications:<br>FC  |  |   |
|                        | Species Specificity:<br>rat   |  |   |
| Background Informatior | Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated $\alpha$ and $\beta$ subunits (PMID: 9779984). CD11b, also known as Integrin alpha M or CR3A, belongs to the integrin alpha chain family. CD11b forms an $\alpha / \beta$ heterodimer with CD18 (integrin $\beta$ 2). CD11b/CD18 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PMID: 9558116; 20008295). CD11b/CD18 is a receptor for the complement protein fragment iC3b, and is also a receptor for fibrinogen, factor X and ICAM1 (PMID: 2971974; 15485828). |  |   |
| Storage                | Storage:<br>Store at 2-8°C. Stable for one year after shipment.<br>Storage Buffer:<br>PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  |  |   |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 LOU rat splenocytes were surface stained with 0.5 ug Anti-Rat CD11b (65229-1-Ig, Clone: OX42) and FITC anti-Mouse IgG2a Antibody at dilution 1:100, or 0.5 ug Control Antibody. Cells were not fixed.