For Research Use Only

Anti-Human CD64 (10.1)

Catalog Number: 65253-1-lg



Purification Method:

Affinity purification

CloneNo.:

10.1

Basic Information

Catalog Number:

65253-1-lg

Size:

100ug, 0.5 mg/mL

Source: Mouse Isotype: IgG1, kappa

Full Name:

P12314

BC032634

2209

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000150337 UNIPROT ID:

Fc fragment of IgG, high affinity Ia,

GenBank Accession Number:

receptor (CD64)

Calculated MW:
374 aa, 43 kDa

Applications

Tested Applications:

FC, ELISA

Species Specificity:

Human

Background Information

Fc γ receptor comprise a multigene family of integral membrane glycoproteins that exhibit complex activation or inhibitory effects on cell functions after aggregation by complexed immunoglobulin G (IgG) (PMID: 17005690). CD64, also known as Fc γ RIA, is a high-affinity receptor for the Fc region of IgG. It is expressed by monocytes/macrophages, activated neutrophils, dendritic cells, and early myeloid cells (PMID: 23293080; 19642859; 7680917). CD64 functions in both innate and adaptive immune responses.

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD64 (65253-1-lg, Clone:10.1) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or Coralite®488-Conjugated Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD64 (65253-1-lg, Clone:10.1) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000, or Coralite®488-Conjugated Mouse IgG1 Isotype Control. Cells were not fixed. Monocytes were gated.