

For Research Use Only

Anti-Human CD11c (3.9) Mouse IgG1 Recombinant Antibody, PBS Only



Catalog Number: 65511-1-PBS

Basic Information

Catalog Number: 65511-1-PBS	GenBank Accession Number: BC038237	Purification Method: Affinity purification
Size: 1mg, 1 mg/ml	GeneID (NCBI): 3687	CloneNo.: 3.9
Source: Mouse	ENSEMBL Gene ID: ENSG00000140678	
Isotype: IgG1	Full Name: integrin, alpha X (complement component 3 receptor 4 subunit)	
	Calculated MW: 1169 aa, 129 kDa	

Applications

Tested Applications:
FC

Species Specificity:
Human

Background Information

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11c, also known as integrin αX , is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α / β heterodimer with CD18 (integrin $\beta 2$). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

Storage

Storage:
Store at -80°C . Stable for one year after shipment.

Storage Buffer:
PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-PBS, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were then stained with FITC-conjugated Anti-Human CD14. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.



1x10⁶ human PBMCs were surface stained with 0.4 ug Anti-Human CD11c Mouse Recombinant Antibody (65511-1-PBS, Clone:3.9) or Mouse IgG1 Isotype Control, and CoraLite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Cells were treated with FC Receptor Block prior to staining. Monocytes were gated.