For Research Use Only

Anti-Human CD21 (BU32) Mouse IgG1 Recombinant Antibody



Purification Method:

Affinity purification

CloneNo.:

BU32

Catalog Number: 65512-1-MR

Basic Information

Catalog Number:

65512-1-MR

Size:

100ug, 0.5 mg/mL

Source: Mouse Isotype: IgG1 GenBank Accession Number:

BC136394 GeneID (NCBI):

1380

ENSEMBL Gene ID:

ENSG00000117322 UNIPROT ID: P20023

Full Name:

complement component (3d/Epstein

Barr virus) receptor 2 Calculated MW: 1092 aa, 119 kDa

Applications

Tested Applications:

FC

Species Specificity:

Human

Background Information

CD21, also known as complement receptor type 2 (CR2), complement C3d receptor, and Epstein-Barr virus receptor, is a transmembrane protein that contains a small cytoplasmic domain, a transmembrane region and an extracellular domain consisting of 15 tandem short consensus repeat sequences (PMID: 6230668; 2551147). It is expressed on B cells, follicular dendritic cells, thymocytes and a subset of peripheral T cells (PMID: 26119182). CD21 binds complement fragments C3d, C3dg and iC3b and acts as a receptor for the Epstein-Barr virus (PMID: 7753047). On B cells, together with CD19 and CD81, CD21 forms a complex that functions as a co-receptor to the BCR (PMID: 7542009). It is involved in B cells activation and functions.

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide.

Selected Validation Data



1x10^6 human PBMCs were surface stained with 0.1 ug Anti-Human CD21 Mouse Recombinant Antibody (65512-1-MR, Clone:BU32) or Mouse IgG1 Isotype Control and Coralite®647-Conjugated AffiniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were then stained with CL488-Conjugated Anti-Human CD3. Cells were not fixed. Lymphocytes were gated.



65512-1-MR CD21 (BU32)

1x10^6 human PBMCs were surface stained with 0.1 ug Anti-Human CD21 Mouse Recombinant Antibody (65512-1-MR, Clone:BU32) or Mouse IgG1 Isotype Control and CoraLite®647-Conjugated AffniPure Donkey Anti-Mouse IgG(H+L) at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.