

For Research Use Only

Anti-Human CD73 (AD2) Mouse IgG1 Recombinant Antibody, PBS Only

Catalog Number: 65531-1-PBS



Basic Information

Catalog Number:

65531-1-PBS

Concentration:

1mg, 2 mg/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

BC015940

GeneID (NCBI):

4907

Full Name:

5'-nucleotidase, ecto (CD73)

Calculated MW:

29 kDa, 63 kDa

Purification Method:

Affinity purification

CloneNo.:

AD2

Applications

Tested Applications:

FC

Species Specificity:

human

Background Information

CD73, also known as ecto-5'-nucleotidase (5'-NT), is a 70-kDa, glycosyl-phosphatidylinositol-linked membrane-bound glycoprotein found in most tissues (PMID: 18404475; 20179192). CD73 is an ectoenzyme that catalyzes the dephosphorylation of AMP and other nucleoside monophosphates (PMID: 9553767). In the human immune system, CD73 is expressed on subsets of T and B cells, on germinal center follicular dendritic cells, and on thymic medullary reticular fibroblasts and epithelial cells (PMID: 2137649; 9553767). CD73 is highly expressed in many human solid tumors and is closely involved in cancer progression (PMID: 20179192).

Storage

Storage:

Store at -80°C. Stable for one year after shipment.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD73 (65531-1-PBS, Clone: AD2) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1) (red), or 0.25 ug Isotype Control and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1) (blue). Cells were not fixed.



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD73 (AD2) Mouse IgG1 RecAb (65531-1-MR, Clone:AD2) or Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21), and PE-conjugated Goat Anti-Mouse IgG. Cells were then stained with APC Anti-Human CD45. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 65531-1-PBS in a different storage



1x10⁶ human PBMCs were surface stained with 0.25 ug Anti-Human CD73 (AD2) Mouse IgG1 RecAb (65531-1-MR, Clone:AD2) (red) or Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue), and PE-conjugated Goat Anti-Mouse IgG. Cells were incubated with FC Receptor Block prior to staining. Cells were not fixed. This data was developed using the same antibody clone with 65531-1-PBS in a different storage buffer formulation.