For Research Use Only

Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody, PBS Only

Catalog Number: 65569-1-PBS



Purification Method:

Protein A purification

CloneNo.:

OKT3

Basic Information

Catalog Number: 65569-1-PBS

GeneID (NCBI): Size:

1mg, 2mg/ml 916 Source: ENSEMBL Gene ID:

Isotype: Full Name:

lgG2a CD3e molecule, epsilon (CD3-TCR complex)

Calculated MW: 207 aa, 23 kDa

ENSG00000198851

BC049847

GenBank Accession Number:

Applications

Tested Applications:

Species Specificity:

human

Mouse

Background Information

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 γ , δ , ϵ , and ζ chains (PMID: 1826255). It is expressed by thy mocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: PMID: PMID3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCRmediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage


Storage:

Store at -80°C. Stable for one year after shipment.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only

Selected Validation Data



1x10^6 human PBMCs were surface stained with Coralite® Plus 647 Anti-Human CD19, and 0.25 ug Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (65569-1-MR, Clone:OKT3) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4), and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65569-1-PBS in a different storage buffer formulation.



1x10^6 human PBMCs were surface stained with 0.25 ug Anti-Human CD3 (OKT3) Mouse IgG2a Recombinant Antibody (65569-1-MR, Clone: OKT3) (red) or 0.25 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue) and PE-Conjugated Goat Anti-Rabbit IgG(H+L). Cells were not fixed. Lymphocytes were gated. This data was developed using the same antibody clone with 65569-1-PBS in a different storage buffer formulation.