

For Research Use Only

Anti-Rat CD80 (3H5) Mouse IgG2a Recombinant Antibody

Catalog Number: 65580-1-MR



Basic Information

Catalog Number:

65580-1-MR

Concentration:

100ug, 1000 ug/ml

Source:

Mouse

Isotype:

IgG2a

GenBank Accession Number:

XM_017597885

GeneID (NCBI):

25408

Full Name:

Cd80 molecule

Calculated MW:

36 kDa

Purification Method:

Protein A purification

CloneNo.:

3H5

Recommended Dilutions:

FC: 0.25 ug per 10⁶ cells in 100 μl suspension

Applications

Tested Applications:

FC

Species Specificity:

rat

Positive Controls:

FC : LPS-stimulated rat splenocytes,

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Storage

Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.09% sodium azide, pH7.3

For technical support and original validation data for this product please contact:



T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ unstimulated or LPS-stimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5) and PE-Conjugated Goat Anti-Mouse IgG. Cells were then stained with CoraLite® Plus 647 Anti-Rat CD45RA. Cells were not fixed.

1x10⁶ LPS-stimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5) (red) or Mouse IgG2a Isotype Control (65208-1-Ig, Clone: C1.18.4) (blue) and PE-Conjugated Goat Anti-Mouse IgG. 1x10⁶ unstimulated rat splenocytes were surface stained with 0.25 ug Anti-Rat CD80/B7-1 Mouse IgG2a Recombinant Antibody (65580-1-MR, Clone: 3H5) (black, dashed) and PE-Conjugated Goat