

For Research Use Only



Anti-Mouse TCR beta (H57-597) Armenian Hamster Recombinant Antibody, PBS Only

Catalog Number: 65678-1-PBS

Basic Information

Catalog Number:

65678-1-PBS

Concentration:

1mg, 3 mg/ml

Source:

Armenian Hamster

Isotype:

IgG2a

GenBank Accession Number:

GeneID (NCBI):

21577

Full Name:

T-cell receptor beta chain

Purification Method:

Protein A purification

CloneNo.:

H57-597

Applications

Tested Applications:

FC

Species Specificity:

mouse

Background Information

T-cell receptor (TCR) consists of heterodimeric glycoproteins (TCR alpha/beta or gamma/delta) that demonstrate homology with immunoglobulins (PMID: 2144901). TCR beta is a member of the immunoglobulin superfamily and is expressed on alpha/beta TCR-bearing T cells and thymocytes. The TCR heterodimers are noncovalently associated with CD3. The TCR/CD3 complex of T-lymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta (PMID: 1826255). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1x10⁶ mouse splenocytes were surface stained with 0.5 ug Anti-Mouse TCR beta Hamster IgG2a RecAb (65678-1-AR, Clone: H57-597) or Armenian Hamster IgG Isotype Control (PIP) (65210-1-Ig, Clone: PIP), and FITC-conjugated Anti-Armenian Hamster IgG antibody. Cells were co-stained with CoraLite® Plus 647 Anti-Mouse CD3. Cells were not fixed. This data was developed using the same antibody clone with 65678-1-PBS in a different storage buffer formulation.



1x10⁶ mouse splenocytes were surface stained with 0.5 ug Anti-Mouse TCR beta Hamster IgG2a RecAb (65678-1-AR, Clone: H57-597) (red) or Armenian Hamster IgG Isotype Control (PIP) (65210-1-Ig, Clone: PIP) (blue), and FITC-conjugated Anti-Armenian Hamster IgG antibody. Cells were not fixed. This data was developed using the same antibody clone with 65678-1-PBS in a different storage buffer formulation.