

RBBP9 Monoclonal antibody

Catalog Number: 66015-2-Ig

Featured Product

Basic Information

Catalog Number:

66015-2-Ig

Size:

1700 µg/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG17864

GenBank Accession Number:

BC015938

GeneID (NCBI):

10741

UNIPROT ID:

O75884

Full Name:

retinoblastoma binding protein 9

Calculated MW:

186 aa, 21 kDa

Observed MW:

22 kDa

Purification Method:

Protein A purification

CloneNo.:

3D5E11

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : pig brain tissue, HeLa cells, A375 cells, A549 cells, rat brain tissue, mouse brain tissue, MCF-7 cells, HepG2 cells, BxPC-3 cells

IHC : human pancreas cancer tissue,

Background Information

RBBP9, also named as BOG, RBBP10, RBBP-9, RBBP-10 and Protein BOG, belongs to the RBBP9 family. It may play a role in the transformation process due to its capacity to confer resistance to the growth-inhibitory effects of TGF- β 1 through interaction with retinoblastoma and the subsequent displacement of E2F-1. RBBP9 is a tumor-associated serine hydrolase activity required for pancreatic neoplasia. It mediates suppression of TGF- β signaling is required for E-cadherin expression as loss of the serine hydrolase activity leads to a reduction in E-cadherin levels and a concomitant decrease in the integrity of tumor cell-cell junctions. RBBP9 protein levels were equivalent in paired primary tumor and nonneoplastic specimens (PMID:20080647)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66015-2-Ig (RBBP9 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66015-2-Ig (RBBP9 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66015-2-Ig (RBBP9 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66015-2-Ig (RBBP9 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



WB result of RBBP9 antibody (66015-2-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RBBP9 transfected A549 cells.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human RBBP9 (66015-2-Ig, Clone:3D5E11) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).