## For Research Use Only

## Bcl-XL Monoclonal antibody, PBS Only (Detector)

proteintech
Antibodies | ELISA kits | Proteins
www.ptglab.com

**Purification Method:** 

Protein A purification

CloneNo.:

4C12A6

Catalog Number: 66020-1-PBS

**Basic Information** 

Catalog Number:

66020-1-PBS

Size:

Source:

Mouse

Isotype:

GenBank Accession Number:

BC019307

GeneID (NCBI):

1mg/ml 59

UNIPROT ID: Q07817 Full Name:

IgG2b BCL2-like 1

Immunogen Catalog Number: Calculated MW: AG18037 26 kDa

Observed MW:

30 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), Cytometric bead array,

Indirect ELISA

Species Specificity:

human, mouse, rat

**Background Information** 

BCL2L1 is a member of the BCL-2 protein family. BCL2L1 is expressed as three isoforms, Bcl-X(L), Bcl-X(s) and Bcl-X(beta), and is located at the outer mitochondrial membrane. The Bcl-XL isoform is a 233 amino acid protein, acting as an apoptotic inhibitor. Bcl-XL can forms heterodimers with BAX, BAK or BCL2, and the heterodimerization with BAX does not seem to be required for anti-apoptotic activity. The Bcl-XS isoform is a shorter variant that is 178 amino acids in length and lacks a 63 amino acid region (amino acids 126-188), acting as an apoptotic activator.

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only


## Selected Validation Data




Jurkat, Ramos, K-562 and Raw 264.7 cells were subjected to SDS PAGE followed by western blot with 66020-1-1g (Bcl-xL Antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66020-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66020-1-lg (Bcl-XL antibody) at dilution of 1:600 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66020-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66020-1-lg (BcL-XL antibody) at dilution of 1:600 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66020-1-PBS in a different storage buffer formulation.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human Bcl-XL (66020-1-1g, Clone:4C12A6) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66020-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Bcl-XL antibody (66020-1-lg, Clone: 4C12A6) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66020-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP50613-1, Bcl-xL Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66020-2-PBS. Detection antibody: 66020-1-PBS. Standard:Ag18037. Range: 1.563-100 ng/mL