

RBAP48 Monoclonal antibody

Catalog Number: 66060-1-Ig

Featured Product

1 Publications

Basic Information

Catalog Number:

66060-1-Ig

Size:

1233 µg/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG6196

GenBank Accession Number:

BC053904

GeneID (NCBI):

5928

UNIPROT ID:

Q09028

Full Name:

retinoblastoma binding protein 4

Calculated MW:

48 kDa

Observed MW:

53 kDa

Purification Method:

Protein A purification

CloneNo.:

5C4D6

Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:20-1:200

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HEK-293 cells, HeLa cells, NIH/3T3 cells, RAW 264.7 cells, Jurkat cell, ROS1728 cells, HSC-T6 cells, L02 cells

IHC : human cervical cancer tissue, human testis tissue

IF/ICC : HepG2 cells,

Background Information

Histone-binding protein RBBP4 (also known as RbAp48, or NURF55) is a protein that in humans is encoded by the RBBP4 gene. This gene encodes a ubiquitously expressed nuclear protein that belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2/NuRD complex complex that has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of corepressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. A decrease of RbAp48 in the dentate gyrus (DG) of the hippocampus in the brain is suspected to be a main cause of memory loss in normal aging (PMID: 23986399).

Notable Publications

Author	Pubmed ID	Journal	Application
Wenxiu Dai	38272027	Dev Cell	

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66060-1-Ig (RBAP48 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 66060-1-Ig (RBAP48 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunohistochemical analysis of paraffin-embedded human cervical cancer using 66060-1-Ig(RBAP48 antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 66060-1-Ig(RBAP48 antibody) at dilution of 1:200 (under 40x lens).



1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug Anti-Human RBAP48 (66060-1-Ig, Clone:5C4D6) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



WB result of RBAP48 antibody (66060-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RBAP48 transfected HeLa cells.