#### For Research Use Only

# TTL Monoclonal antibody

Catalog Number:66076-1-lg

Featured Product

7 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:
66076-1-lg BC036819
Concentration: GeneID (NCBI):
900 ug/ml 150465
Source: UNIPROT ID:
Mouse Q8NG68

IgG2b tubulin tyrosine ligase

Immunogen Catalog Number:Calculated MW:AG4708377 aa, 43 kDa

Observed MW: 43 kDa

Full Name:

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB. IF

Isotype:

Species Specificity: human, mouse, rat, pig

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: LNCaP cells, fetal human brain tissue, SH-SY5Y cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:20-1:200

CloneNo.:

2E5F8

IHC: human liver tissue,

## **Background Information**

Tubulin-tyrosine ligase (TTL) is the enzyme responsible for the reversible addition of a tyrosine residue at the carboxyl end of alpha-tubulin. TTL forms stable complexes with tubulin and inhibit tubulin polymerization. TTL is frequently suppressed during tumor progression with resulting accumulation of detyrosinated alpha-tubulin in tumor cells. TTL suppression in human cancers is associated with increased tumor aggressiveness.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Luísa T Ferreira	29804676	Methods Cell Biol	
Amrendra Mishra	31170286	Carcinogenesis	
Rui Wang	31408157	J Mol Cell Biol	IF

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


### Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66076-1-1g (TTL antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver using 66076-1-lg(TTL antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver using 66076-1-1g(TTL antibody) at dilution of 1:50 (under 40x lens).