For Research Use Only

PDCD4 Monoclonal antibody

Catalog Number:66100-1-lg Featured Product

3 Publications



Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:1000-1:8000 IHC 1:50-1:500

CloneNo.:

1F1A11

Basic Information

Catalog Number: 66100-1-lg Size:

1400 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG19239

GenBank Accession Number: BC026104

GeneID (NCBI): 27250 **UNIPROT ID:** Q53EL6

programmed cell death 4 (neoplastic transformation inhibitor)

Calculated MW: 469 aa. 52 kDa Observed MW: 56 kDa

Full Name:

Applications

Tested Applications: IHC, WB, ELISA **Cited Applications:**

Species Specificity: human **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK293 cells IHC: human liver cancer tissue,

Background Information

 $Programmed \ cell\ death\ 4\ (Pdcd4)\ is\ a\ novel\ tumor\ suppressor\ that\ inhibits\ translation,\ progression\ and\ invasion.\ It$ was first identified as being differnetially upregulated during apoptosis. Pdcd4 interferes with the activity of the eukaryotic initiation factor (eIF) 4A by displacing the scaffold protein eIF4G from its binding to the RNA helicase

Notable Publications

Author	Pubmed ID	Journal	Application
Ding Jiamin	36083475	J Tradit Chin Med	WB
Ting Peng	35695407	J Med Chem	WB
Changyu Chen	30522853	Clin Res Hepatol Gastroenterol	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of PDCD4 antibody (66100-1-1g, 1:4000) with si-control and si-PDCD4 transfected HEK293 cells.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66100-1-1g (PDCD4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HeLa cells were subjected to SDS PAGE followed by western blot with 66100-1-1g (PDCD4 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66100-1-1g (PDCD4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).