

For Research Use Only

# FUT8 Monoclonal antibody

Catalog Number: 66118-1-Ig

Featured Product

13 Publications



## Basic Information

Catalog Number:

66118-1-Ig

Size:

2300 µg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG18769

GenBank Accession Number:

BC025385

GeneID (NCBI):

2530

UNIPROT ID:

Q9BYC5

Full Name:

fucosyltransferase 8 (alpha (1,6)

fucosyltransferase)

Calculated MW:

66 kDa

Observed MW:

66 kDa

Purification Method:

Protein A purification

CloneNo.:

2F7H2

Recommended Dilutions:

WB 1:1000-1:8000

IHC 1:250-1:1000

## Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human

Cited Species:

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

**WB** : MDA-MB-231 cells, MDA-MB-453s cells, MCF-7 cells, Caco-2 cells

**IHC** : human colon cancer tissue, human breast cancer tissue, human lung cancer tissue

## Background Information

Fucosyltransferase 8 (FUT8) is GDP-fucose-glycoprotein fucosyltransferase and is localized in Golgi apparatus. FUT8 is involved in protein glycosylation by catalyzing the addition of fucose in alpha 1-6 linkage to the first GlcNAc residue, next to the peptide chains in N-glycans. FUT8 may be involved in human aging by regulating protein N-glycosylation and in the molecular pathogenesis of human pulmonary emphysema (PE).

## Notable Publications

Author	Pubmed ID	Journal	Application
Toni M Mueller	27773385	Schizophr Res	
López-Cortés Rubén	33021632	Glycobiology	WB
Paulina Sosicka	36053214	J Cell Biol	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




MDA-MB-231 cells were subjected to SDS PAGE followed by western blot with 66118-1-Ig (FUT8 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of FUT8 antibody (66118-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FUT8 transfected MCF-7 cells.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66118-1-Ig (FUT8 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66118-1-Ig (FUT8 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).