### For Research Use Only

# IL-9 Monoclonal antibody

Catalog Number:66144-1-lg 5 Publications



**Basic Information** 

Catalog Number: 66144-1-lg Size:

2400 ug/ml

Source: Mouse

Isotype:

lgG2b

Immunogen Catalog Number:Calculated MW:AG21435144 aa, 16 kDa

Observed MW: 19 kDa

BC066284

3578

P15248

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

interleukin 9

GenBank Accession Number:

Purification Method:

Protein A purification

CloneNo.: 1F3A11

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:200-1:800 IF-P 1:200-1:800

**Applications** 

Tested Applications: WB, IHC, IF-P, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:** 

WB: Recombinant protein, Recombinant protein

protein

IHC: human tonsillitis tissue,
IF-P: human tonsillitis tissue,

# **Background Information**

Interleukin 9 (IL-9) is a  $\gamma$  c-family cytokine that is highly produced by T-helper 9 (Th9) cells. IL-9 could promote the survival and activation of various cellular targets, including mast cells, B cells, T cells, and structural cells. Its expression is considered a hallmark of Th2-lineage cells. Primarily studied in Th2-type immunity, IL-9 was shown to be involved in asthma, allergy, and host defense against helminth infections. IL-9 also stimulates cell proliferation and prevents apoptosis. It functions through the interleukin 9 receptor (IL9R), which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yujun Sheng	33249782	FEBS Open Bio	WB
Tanfang Zhou	39199394	Biomolecules	IHC,IF
Liang Yan	37641663	Clin Cosmet Investig Dermatol	IHC

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Recombinant protein were subjected to SDS PAGE followed by western blot with 66144-1-1g (II.9 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using IL-9 antibody (66144-1-lg, Clone: 1F3A11) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66144-1-1g (IL-9 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66144-1-1g (IL-9 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).