## For Research Use Only

## IL-6 Monoclonal antibody, PBS Only

Catalog Number: 66146-1-PBS

**Featured Product** 



**Basic Information** 

Catalog Number:

GenBank Accession Number:

**Purification Method:** Protein A purification

66146-1-PBS

Size:

GeneID (NCBI): 3569

BC015511

CloneNo.: 1A3B4

1 mg/ml Source:

ENSEMBL Gene ID: ENSG00000136244

Mouse Isotype:

**UNIPROT ID:** P05231

Immunogen Catalog Number: AG10646

lgG2b

Full Name:

interleukin 6 (interferon, beta 2)

Calculated MW: 212 aa, 24 kDa Observed MW: 24 kDa

**Applications** 

**Tested Applications:** 

WB,ELISA

Species Specificity:

human

**Background Information** 

Interleukin-6 (IL-6) is an interleukin that acts as both a pro-inflammatory and anti-inflammatory cytokine. IL-6protein is secreted by a variety of cell types including T cells and macrophages as phosphorylated and variably glycosylated molecule. IL-6 plays an essential role in the final differentiation of B-cells into Ig-secreting cells involved in lymphocyte and monocyte differentiation. It induces myeloma and plasmacytoma growth and induces nerve cells differentiation Acts on B-cells, T-cells, hepatocytes, hematopoietic progenitor cells and cells of the CNS. IL-6 is also considered a myokine, a cytokine produced from muscle, and is elevated in response to muscle contraction. IL-6 has been shown to interact with interleukin-6 receptor and glycoprotein 130. Additionally, IL-6 is involved in hematopoiesis, bone metabolism, and cancer progression, and has been defined an essential role in directing transition from innate to acquired immunity.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Non-treated and protein transport inhibitor treated U-87 MG cells were subjected to SDS PAGE followed by western blot with 66146-1-lg (IL-6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66146-1-PBS in a different storage buffer formulation.



Non-treated, LPS and protein transport inhibitor treated HUVEC cells were subjected to SDS PAGE followed by western blot with 66146-1-lg (IL-6 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66146-1-PBS in a different storage buffer formulation.