For Research Use Only

IFN Alpha Monoclonal antibody

Catalog Number:66162-1-lg <u>3 Publications</u>




Basic Information	Catalog Number: 66162-1-lg	GenBank Accession Number: BC074928	Purification Method: Protein A purification
	Size: 2700 µg/ml	GeneID (NCBI): 3439	CloneNo.: 4A2A11
	Source: Mouse	UNIPROT ID: P01562	Recommended Dilutions: WB 1:500-1:1000
	Isotype: IgG2a	Full Name: IFNA1	
	Immunogen Catalog Number: AG12533	Calculated MW: 189 aa, 22 kDa	
		Observed MW: 22 kDa	
Applications	Tested Applications: WB, ELISA	Positive Controls:	
	Cited Applications: WB	WB : Recombinant protein, Transfected HEK-293 cells	
	Species Specificity: human		
	Cited Species: human		
Background Information	IFNA1 (IF-alpha) is a member of the Type I IFN (1) family best known for their antiviral activity. It is a key cytokine regulating the activity of B cells, T-helper cells (Th cells), cDCs and natural killer cells (NK cells). IFNA1 induces B cell maturation into plasma cells and immunoglobulin production. IFNA1 plays an important role in the pathogenesis of systemic lupus erythematosus (SLE). IFNA1 was the first cytokine to show clinical benefit in the treatment of certain types of cancer, including melanoma, chronic myelogenous leukemia, and renal cancer.		
Notable Publications	Author	Pubmed ID Journal	Application
	Yan Jin	34524541 Amino Acids	WB
	Jianguo Feng	31211948 Brain Res	WB
	Juanjuan Luo	37027305 Cell Rep	WB
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



WB result of Interferon alpha antibody (66162-1-lg; 1:300; room temperature for 1.5 hours) with negative control and Interferon alpha overexpressed Transfected HEK-293 cells.

Recombinant protein were subjected to SDS PAGE followed by western blot with 66162-1-1g (Interferon alpha antibody at dilution of 1:40000 incubated at room temperature for 1.5 hours.