For Research Use Only

MSH6 Monoclonal antibody

Catalog Number:66172-1-lg Featured Product

5 Publications



Basic Information

Catalog Number: GenBank Accession Number: 66172-1-lg BC004246 GeneID (NCBI): Size: 1000 µg/ml 2956 **UNIPROT ID:** Source:

Mouse P52701 Full Name: Isotype:

IgG2a mutS homolog 6 (E. coli) Calculated MW: Immunogen Catalog Number:

AG12645 153 kDa

> Observed MW: 160 kDa

Purification Method: Protein A purification

CloneNo.: 2E10B2

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:500-1:2000 IF 1:50-1:500

Applications

Tested Applications: FC, IF, IHC, IP, WB, ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse **Cited Species:**

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HepG2 cells, U2OS cells, A431 cells, HeLa cells, MCF-7 cells, K-562 cells

IP: HEK-293 cells,

IHC: human colon cancer tissue,

IF: HepG2 cells,

Background Information

MSH6, also named as DNA mismatch repair protein Msh6 or GTMBP, is a 1360 amino acid protein, which contains one PWWP domain and belongs to the DNA mismatch repair MutS family. MSH6 localizes in the nucleus and is a component of the post-replicative DNA mismatch repair system. Msh2 and Msh6 form a protein complex required to repair mismatches generated during DNA replication.

Notable Publications

Author	Pubmed ID	Journal	Application
Takashi Minamisaka	35614379	Neuropathology	IHC
Jun Zhu	32396667	J Surg Oncol	IHC
Dingkong Liang	29352574	Surgery	IHC

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66172-1-lg (MSH6 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1X10^6 HEK-293 cells were intracellularly stained with 0.2 ug Anti-Human MSH6 (66172-1-1g, Clone:2E10B2) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (black). Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100.




WB result of MSH6 antibody (66172-1-lg, 1:1000) with si-Control and si-MSH6 transfected HEK293 cells



HEK-293 cells were subjected to SDS PAGE followed by western blot with 66172-1-1g (MSH6 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



IP result of anti-MSH6 (IP:66172-1-Ig, 4ug; Detection:66172-1-Ig 1:600) with HEK-293 cells lysate 1800ug.



Immunofluorescent analysis of (10% Formaldehyde) fixed HepG2 cells using 66172-1-1g(MSH6 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+I).

U2OS cells were subjected to SDS PAGE followed by western blot with 66172-1-1g (MSH6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Various lysates were subjected to SDS PAGE followed by western blot with 66172-1-Ig (MSH6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.