For Research Use Only

MSH6 Monoclonal antibody

Catalog Number:66172-1-lg

Featured Product

9 Publications



Basic Information

 Catalog Number:
 GenBank Accession Number:

 66172-1-lg
 BC004246

 Size:
 GeneID (NCBI):

 1000 ug/ml
 2956

 Source:
 UNIPROT ID:

 Mouse
 P52701

Nouse P52/01
Isotype: Full Name:
IgG2a mutS homolog 6 (E. coli)

Immunogen Catalog Number: Calculated MW:

AG12645 153 kDa

Observed MW: 160 kDa

Positive Controls:

WB: HEK-293 cells, HepG2 cells, U2OS cells, A431 cells, HeLa cells, MCF-7 cells, K-562 cells

Purification Method:

Protein A purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

WB 1:500-1:2000

IHC 1:500-1:2000

IF/ICC 1:500-1:2000

protein lysate

CloneNo.:

2E10B2

IP: HEK-293 cells,

IHC: human colon cancer tissue, Ramos cells

IF/ICC: U2OS cells,

Applications

Tested Applications: WB, IHC, IF/ICC, IP, ELISA Cited Applications:

WB, IHC

Species Specificity: human, mouse Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

MSH6, also named as DNA mismatch repair protein Msh6 or GTMBP, is a 1360 amino acid protein, which contains one PWWP domain and belongs to the DNA mismatch repair MutS family. MSH6 localizes in the nucleus and is a component of the post-replicative DNA mismatch repair system. Msh2 and Msh6 form a protein complex required to repair mismatches generated during DNA replication.

Notable Publications

Author	Pubmed ID	Journal	Application
Takashi Minamisaka	35614379	Neuropathology	IHC
Jun Zhu	32396667	J Surg Oncol	IHC
Dingkong Liang	29352574	Surgery	IHC

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66172-1-lg (MSH6 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed U2OS cells using MSH6 antibody (66172-1-Ig, Clone: 2E10B2) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).




WB result of MSH6 antibody (66172-1-lg, 1:1000) with si-Control and si-MSH6 transfected HEK293




HEK-293 cells were subjected to SDS PAGE followed by western blot with 66172-1-1g (MSH6 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




IP result of anti-MSH6 (IP:66172-1-Ig, 4ug; Detection:66172-1-Ig 1:600) with HEK-293 cells lysate 1800ug.




U2OS cells were subjected to SDS PAGE followed by western blot with 66172-1-lg (MSH6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66172-1-1g (MSH6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.





1x10^6 HEK-293 cells were intracellularly stained with 0.4 ug MSH6 Monoclonal antibody (66172-1-lg, Clone:2E10B2) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1) (red), or 0.4 ug Mouse IgG2a isotype control Mouse McAb (66360-2-lg, Clone: 11A1B2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).