

Vitamin D binding protein

Monoclonal antibody

 Catalog Number: **66175-1-Ig** **1 Publications**

Basic Information

Catalog Number: 66175-1-Ig	GenBank Accession Number: BC057228	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 2638	CloneNo.: 1E4D10
Source: Mouse	UNIPROT ID: P02774	Recommended Dilutions: WB 1:1000-1:8000 IHC 1:250-1:1000
Isotype: IgG1	Full Name: group-specific component (vitamin D binding protein)	
Immunogen Catalog Number: AG9803	Calculated MW: 474 aa, 53 kDa	
	Observed MW: 52-58 kDa	

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA	Positive Controls:
Cited Applications: WB, IHC, CoIP, ChIP, IF	WB : human testis tissue,
Species Specificity: human	IHC : human liver tissue, human colon cancer tissue, human liver cancer tissue
Cited Species: mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

Vitamin D binding protein is a sparsely glycosylated serum protein responsible for highly specific binding and tissue-specific delivery of vitamin D and its metabolites. In addition, it is also an actin scavenger, and is the precursor to the immunomodulatory protein, Gc-MAF. Vitamin D binding protein has been proposed to have significant roles in C5a chemotaxis, osteoclast development and possibly in macrophage activation/recruitment.

Notable Publications

Author	Pubmed ID	Journal	Application
Lu-Ning Qin	38164156	Theranostics	IHC, IF, WB, CoIP, ChIP

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human testis tissue were subjected to SDS PAGE followed by western blot with 66175-1-Ig (Vitamin D binding protein antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66175-1-Ig (Vitamin D binding protein antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66175-1-Ig (Vitamin D binding protein antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66175-1-Ig (Vitamin D binding protein antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ U-937 cells were intracellularly stained with 0.4 ug Anti-Human Vitamin D binding protein (66175-1-Ig, Clone:1E4D10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).