For Research Use Only

SMARCE1 Monoclonal antibody

Catalog Number:66182-1-Ig



Purification Method:

CloneNo.: 1F3D6

IHC 1:20-1:200

IF/ICC 1:200-1:800

Protein G purification

Recommended Dilutions: WB 1:500-1:2000

Basic Information

Catalog Number: 66182-1-lg Size: 1800 ug/ml

Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG1161

GenBank Accession Number:

BC007082 GeneID (NCBI): 6605

UNIPROT ID:

Q969G3 **Full Name:**

SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily e, member 1

Calculated MW: 47 kDa Observed MW: 50-55 kDa

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval antigen retrieval with a trace buffer and 16.0 with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells,

IHC: human breast cancer tissue.

IF/ICC: HEK-293 cells.

Background Information

 $Via\ antagonizing\ chromatin-mediated\ transcriptional\ repression, the\ SWI/SNF\ complex\ in\ S.\ cerevisiae\ and\ solve the support of th$ $Drosophila\ is\ thought\ to\ facilitate\ transcriptional\ activation\ of\ specific\ genes.\ The\ BRG1/brm-associated\ factors,\ or\ activation\ of\ specific\ genes.$ SMARCE1, complex in mammals is functionally related to SWI/SNF and consists of 9 to 12 subunits, some of which are homologous to SWI/SNF subunits. It contains a DNA-binding HMG domain and performs multiple biological functions, including DNA-, protein- and chromatin-binding, transcription coactivating activity, chromatin modification, DNA-dependent negative regulation of transcription, regulation of transcription of the RNA polymerase II promoter. In addition, SMARCE1, as an ER (estrogen receptor) subtype-selective modulator, regulates ER mediated transcription in a ligand-dependent manner and also directly binds to AR (androgen receptor) to alter its activity. The calculated molecular weight of SMARCE1 is 47 kDa, but modified SMARCE1 is about 55 kDa.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 66182-1-1g (SMARCE1 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66182-1-1g (SMARCE1 Antibody) at dilution of 1:100 (under 40x lens).



Immunofluorescent analysis of HEK-293 cells using 66182-1-Ig (SMARCE1 antibody) at dilution of 1:50 and FITC-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using SMARCE1 antibody (66182-1-Ig, Clone: 1F3D6) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002), CL594-Phalloidin (red).