For Research Use Only

## PD-1/CD279 Monoclonal antibody, PBS Only

www.ptglab.com

**Purification Method:** 

CloneNo.:

4H4D1

Protein A purification

Catalog Number: 66220-1-PBS

Featured Product

**Basic Information** 

Catalog Number:

66220-1-PBS

Size: 1000 µg/ml

Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG12470

programmed cell death 1 Calculated MW: 288 aa, 32 kDa

BC074740

GeneID (NCBI):

**UNIPROT ID:** 

Q15116 Full Name:

GenBank Accession Number:

Observed MW: 32 kDa, 47-55 kDa

**Applications** 

**Tested Applications:** Species Specificity: human, mouse, rat, pig

WB, IHC, IF-P, ELISA

## **Background Information**

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one lg superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytolytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436). The calculated molecular weight of PD-1 is 32 kDa. It has been reported that PD-1 is heavily glycosylated and migrates with an apparent molecular mass of 47-55 kDa on SDS-PAGE (PMID: 8671665; 17640856; 17003438).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66220-1-1g (PD-1/CD279 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 66220-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66220-1-Ig (PD-1/CD279 antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66220-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using PD-1/CD279 mouse mAb (66220-1-Ig) at dilution of 1:50 and CD20 rabbit pAb (24828-1-AP) at dilution of 1:50, further stained with Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L) for 66220-1-Ig, and Alexa Fluor 594-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) for 24828-1-AP. This data was developed using the same antibody clone with 66220-1-PBS in a different storage buffer



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using PD-1/CD279 antibody (66220-1-lg, Clone: 4H4D1) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L). This data was developed using the same antibody clone with 66220-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66220-1-1g (PD-1/CD279 antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66220-1-PBS in a different storage buffer formulation.