For Research Use Only

Beta Tubulin Monoclonal antibody, PBS Only

www.ptglab.com

Catalog Number: 66240-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

1D4A4

66240-1-PBS

GeneID (NCBI):

CloneNo.:

Size: 1mg/ml Source:

10381

BC000748

UNIPROT ID:

Mouse Isotype: Q13509 Full Name: tubulin, beta 3

lgG2a Immunogen Catalog Number:

Calculated MW:

AG0117

450 aa, 50 kDa

Observed MW: 50-55 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

Species Specificity:

human, mouse, rat, pig, zebrafish, nematode

Background Information

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart, once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha and beta tubulins have been widely used as loading controls. Tubulin expression may vary according to resistance to antimicrobial and antimitotic drugs.

Storage


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66240-1-lg (beta Tubulin antibody) at dilution of 1:200000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation.





Various lysates were subjected to SDS PAGE followed by western blot with 66240-1-1g (beta Tubulin antibody) at dilution of 1:100000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 66240-1-lg (beta Tubulin antibody), at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Red: staning with Coralite555-Phalloidin. This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation.








IP result of anti-Beta Tubulin (IP:66240-1-Ig, 5ug; Detection:66240-1-Ig 1:1000) with HeLa cells lysate 880ug. This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse eye tissue using Beta Tubulin antibody (66240-1-lg, Clone: 1D4A4) at dilution of 1:800 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using Beta Tubulin antibody (66240-1-lg, Clone: 1D4A4) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 66240-1-PBS in a different storage buffer formulation