For Research Use Only

CD1d Monoclonal antibody

Catalog Number:66257-1-lg Featured Product

1 Publications



Purification Method:

CloneNo.:

1C12F1

Positive Controls:

WB: human testis tissue, Jurkat

Protein G purification

Recommended Dilutions:

IHC: human skin tissue, human small intestine tissue

WB 1:500-1:2000

IHC 1:20-1:200

Basic Information

Catalog Number: 66257-1-lg

Size:

GenBank Accession Number:

GeneID (NCBI): 912

2000 μg/ml Source: **UNIPROT ID:** P15813 Mouse Isotype: Full Name: CD1d molecule

Calculated MW: Immunogen Catalog Number: AG11108 335 aa, 38 kDa

> Observed MW: 40 kDa, 50-55 kDa

Applications

Tested Applications:

IHC, WB,ELISA

Cited Applications:

WB. IF

Species Specificity:

human

Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval antigen retrieval antigen retrieval and 6.0 with citrate buffer pH 6.0

Background Information

 ${\tt CD1d}\ is\ a\ member\ of\ the\ CD1\ family\ of\ transmembrane\ glycoproteins, which\ are\ structurally\ related\ to\ the\ major\ properties and the constraints of\ the\ con$ histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. CD1d is an antigenpresenting protein that binds self and non-self glycolipids and presents them to T-cell receptors on NKT cells. When activated, NKT cells rapidly produce Th1 and Th2 cytokines. The molecular weight of unglycosylated CD1d is 38 kDa, while glycosylated form of CD1d is 48-55 kDa.

Notable Publications

Author Pubmed ID Journal Application WB.IF 38868316 Qingyuan Liu Open Med (Wars)


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of CD1D antibody (66257-1-lg, 1:500) with si-Control and si-CD1D transfected Jurkat cells.



human testis tissue were subjected to SDS PAGE followed by western blot with 66257-1-1g (CD1D Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human skin tissue slide using 66257-1 Ig (CD1d antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human skin tissue slide using 66257-1g (CD1d antibody) at dilution of 1:100 (under 40x lens).