

For Research Use Only

# KGA-Specific Monoclonal antibody, PBS Only



Catalog Number: 66265-2-PBS

## Basic Information

Catalog Number: 66265-2-PBS	GenBank Accession Number: BC038507	Purification Method: Protein G purification
Size: 1 mg/ml	GeneID (NCBI): 2744	CloneNo.: 1B11A11
Source: Mouse	UNIPROT ID: O94925	
Isotype: IgG1	Full Name: glutaminase	
Immunogen Catalog Number: AG24227	Calculated MW: 669 aa, 73 kDa	
	Observed MW: 65 kDa	

## Applications

Tested Applications:  
WB, Indirect ELISA, IHC, IF

Species Specificity:  
human, mouse, rat

## Background Information

GLS, also named as GLS1 and KIAA0838, belongs to the glutaminase family. It catalyzes the first reaction in the primary pathway for the renal catabolism of glutamine. Glutaminase-, glutamate-, and taurine-immunoreactive neurons develop neurofibrillary tangles in Alzheimer's disease. The glutaminase band in AA/C1 cells is more intense than in HT29 cells, in accordance with measurements of glutaminase activity, and had the same molecular mass of approx. 63 kDa. The bands for both cell lines are clearly different in size from both rat liver glutaminase (58 kDa) and rat kidney glutaminase (65 kDa) (PMID: 12408749). It also reveals a molecular weight of 83-84 kDa as a phosphate-dependent glutaminase (PMID: 447624; 7512428). It has 3 isoforms produced by alternative splicing named as KGA, GAM, GAC. This antibody is specific to KGA.

## Storage


Storage:  
Store at -80°C.  
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:  
PBS Only


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 66265-2-Ig (GLS antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66265-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 66265-2-Ig (GLS antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66265-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 66265-2-Ig (GLS antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66265-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using KGA-Specific antibody (66265-2-Ig, Clone: 1B11A11) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66265-2-PBS in a different storage buffer formulation.