## For Research Use Only

## Gelsolin Monoclonal antibody

Catalog Number:66280-1-lg Featured Product



**Basic Information** 

Catalog Number: GenBank Accession Number: 66280-1-lg BC026033

66280-1-lg BC026033 Thiophilic affinity chromatograph
Concentration: GeneID (NCBI): CloneNo.:
531 ug/ml 2934 3C4A2

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Mouse
 P06396
 WB 1:1000-1:4000

 Isotype:
 Full Name:
 IHC 1:20-1:1000

IgA gelsolin (amyloidosis, Finnish type)

Immunogen Catalog Number: Calculated MW: 782 aa, 86 kDa
Observed MW:

80-83 kDa

**Applications** 

Tested Applications: WB, IHC, ELISA Species Specificity:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

human

Positive Controls:

WB: human plasma tissue, U2OS cells, A549 cells,

**Purification Method:** 

human plasma, HT-29 cells

IHC: human colon tissue, human colon cancer tissue

## **Background Information**

Gelsolin (GSN) is an actin binding protein that regulates actin-severing/depolymerizing in a calcium dependent manner, and acts as a multifunctional regulator for cell metabolism and survival. Gelsolin exists in two isoforms: a cytoplasmic gelsolin (cGSN; 80-90 kDa) and a plasma gelsolin (pGSN; 93 kDa), differing at N terminal residues. Decreased levels of pGSN are strongly associated with human diseases, such as, acute liver injury, major trauma, myocardial infarction, inflammation, lung injury, rheumatoid arthritis, hemodialysis, multiple sclerosis, Alzheimer's disease.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




U2OS (WT and GSN KO) lysates prepared with RIPA buffer, 25 µg protein loaded, 66280-1-1g incubated at 1:1000 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency.



human plasma were subjected to SDS PAGE followed by western blot with 66280-1-1g (Gelsolin Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66280-1-Ig (Gelsolin Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66280-1-Ig (Gelsolin Antibody) at dilution of 1:200 (under 40x lens).