

For Research Use Only

FABP5 Monoclonal antibody

Catalog Number: 66299-1-Ig

Featured Product

3 Publications



Basic Information

Catalog Number:

66299-1-Ig

Size:

1400 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG3005

GenBank Accession Number:

BC019385

GeneID (NCBI):

2171

UNIPROT ID:

Q01469

Full Name:

fatty acid binding protein 5 (psoriasis-associated)

Calculated MW:

135 aa, 15 kDa

Observed MW:

15 kDa

Purification Method:

Protein A purification

CloneNo.:

1C6E12

Recommended Dilutions:

WB 1:2000-1:16000

IHC 1:200-1:4000

IF 1:200-1:800

Applications

Tested Applications:

FC, IF/ICC, IHC, WB, ELISA

Cited Applications:

WB, IHC

Species Specificity:

human, mouse, rat

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A375 cells, U2OS cells, rat brain tissue, fetal human brain tissue, A549 cells, HeLa cells, HEK-293 cells, mouse brain tissue

IHC: human breast cancer tissue, human prostate cancer tissue, mouse brown adipose tissue

IF: HepG2 cells,

Background Information

FABP5, also named as PA-FABP and E-FABP, belongs to the calycin superfamily and Fatty-acid binding protein (FABP) family. It is high specificity for fatty acids. FABP5 is highest affinity for C18 chain length. It may be involved in keratinocyte differentiation. FABP5 is a fatty acid-binding protein and is expressed in epidermis and endothelial cells of the microvasculature of different organs. FABP5 has also been identified as a tumor-associated antigen, which is highly expressed in various cancers. FABP5 was detected in the sera of HNSCC patients with early stage cancer. Antibodies specific for FABP5 were significantly increased in a substantial amount in patients, suggesting that FABP5 may be a potential diagnostic biomarker for HNSCC. FABP5 may serve as a biomarker for HNSCC. (PMID:19602232)

Notable Publications

| Author | Pubmed ID | Journal | Application |
|----------------|-----------|-----------------|-------------|
| Masafumi Ohira | 33754641 | Carcinogenesis | WB,IHC |
| Jinghui Lu | 34976793 | Front Oncol | WB |
| Risa Nakagawa | 31432248 | Med Mol Morphol | IHC |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of FABP5 antibody (66299-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FABP5 transfected A375 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using FABP5 antibody (66299-1-Ig, Clone: 1C6E12) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1X10⁶ HeLa cells were intracellularly stained with 0.20 ug/test Anti-Human FABP5 (66299-1-Ig, Clone:1C6E12) (red) or 0.20 ug control antibody (blue) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) with dilution 1:1000. Fixed with 90% MeOH.






A375 cells were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 Antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).