

FABP5 Monoclonal antibody

Catalog Number: 66299-1-Ig

Featured Product

5 Publications

Basic Information

Catalog Number: 66299-1-Ig	GenBank Accession Number: BC019385	Purification Method: Protein G purification
Source: Mouse	GeneID (NCBI): 2171	CloneNo.: 1C6E12
Isotype: IgG1	UNIPROT ID: Q01469	Recommended Dilutions: WB: 1:2000-1:16000 IHC: 1:200-1:4000 IF/ICC: 1:200-1:800
Immunogen Catalog Number: AG3005	Full Name: fatty acid binding protein 5 (psoriasis-associated)	
	Calculated MW: 135 aa, 15 kDa	
	Observed MW: 15 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, ELISA

Cited Applications:
WB, IHC, IF

Species Specificity:
human, mouse, rat

Cited Species:
human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A375 cells, fetal human brain tissue, Hepa1-6 cells, rat brain tissue, U2OS cells, mouse brain tissue, A549 cells, HeLa cells, HEK-293 cells

IHC: human breast cancer tissue, human prostate cancer tissue, mouse brown adipose tissue

IF/ICC: HepG2 cells,

Background Information

FABP5, also named as PA-FABP and E-FABP, belongs to the calycin superfamily and Fatty-acid binding protein (FABP) family. It is high specificity for fatty acids. FABP5 is highest affinity for C18 chain length. It may be involved in keratinocyte differentiation. FABP5 is a fatty acid-binding protein and is expressed in epidermis and endothelial cells of the microvasculature of different organs. FABP5 has also been identified as a tumor-associated antigen, which is highly expressed in various cancers. FABP5 was detected in the sera of HNSCC patients with early stage cancer. Antibodies specific for FABP5 were significantly increased in a substantial amount in patients, suggesting that FABP5 may be a potential diagnostic biomarker for HNSCC. FABP5 may serve as a biomarker for HNSCC. (PMID:19602232)

Notable Publications

Author	Pubmed ID	Journal	Application
Masafumi Ohira	33754641	Carcinogenesis	WB, IHC
Jinghui Lu	34976793	Front Oncol	WB
Risa Nakagawa	31432248	Med Mol Morphol	IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
 Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




WB result of FABP5 antibody (66299-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FABP5 transfected A375 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




WB result of FABP5 antibody (66299-1-Ig; 1:1000; room temperature for 1.5 hours) with wild-type and FABP5 knockout Hepa1-6 cells.




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using FABP5 antibody (66299-1-Ig, Clone: 1C6E12) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunohistochemical analysis of paraffin-embedded mouse brown adipose tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).






A375 cells were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 Antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66299-1-Ig (FABP5 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66299-1-Ig (FABP5 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 66299-1-Ig (FABP5 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).