

CD27 Monoclonal antibody

Catalog Number: 66308-1-Ig **1 Publications**

Basic Information

Catalog Number: 66308-1-Ig	GenBank Accession Number: BC012160	Purification Method: Protein G purification
Concentration: 1500 ug/ml	GeneID (NCBI): 939	CloneNo.: 1C1G3
Source: Mouse	ENSEMBL Gene ID: ENSG00000139193	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000 IF-P 1:200-1:800
Isotype: IgG1	UNIPROT ID: P26842	
Immunogen Catalog Number: AG24537	Full Name: CD27 molecule	
	Calculated MW: 29 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: WB, IHC, IF-P, FC, ELISA	Positive Controls: WB : Raji cells, pig spleen tissue, Ramos cells, Daudi cells IHC : human tonsillitis tissue, IF-P : human tonsillitis tissue, FC : human PBMCs,
Cited Applications: IF	
Species Specificity: human, pig	
Cited Species: human	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

CD27 (also known as TNFRSF7) is a type I glycoprotein expressed on some B cells and the majority of T cells, and is a member of the tumor necrosis factor (TNF) receptor family. CD27 is required for generation and long-term maintenance of T cell immunity (PMID: 11062504). It is a receptor for CD70 (CD27L). Ligand of CD27 by CD70 induces strong ubiquitination of TRAF and the activation of both canonical and non-canonical NF-kappaB pathways, as well as the JNK pathway (PMID: 19426224). CD27 may also play a role in apoptosis through association with SIVA1.

Notable Publications

Author	Pubmed ID	Journal	Application
Yang Liu	39231979	Nat Commun	IF

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66308-1-Ig (CD27 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66308-1-Ig (CD27 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66308-1-Ig (CD27 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66308-1-Ig (CD27 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1x10⁶ human PBMCs were surface stained with 0.2 μ g CD27 Monoclonal antibody (66308-1-Ig, Clone:1C1G3) or 0.2 μ g Mouse IgG1 isotype control (66360-1-Ig, Clone: T1F8D3F10), and CoraLite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO.RGAM005). Cells were not fixed.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66308-1-Ig (CD27 antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using CD27 antibody (66308-1-Ig, Clone: 1C1G3) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).