For Research Use Only

Peripherin Monoclonal antibody

Catalog Number: 66317-1-lg



Basic Information

Catalog Number: 66317-1-lg

Concentration:

1000 ug/ml

Source:

BC032703 GeneID (NCBI): 5630

GenBank Accession Number:

UNIPROT ID: Mouse P41219 Full Name: Isotype: lgG2b peripherin Calculated MW: Immunogen Catalog Number: AG11116 470 aa, 54 kDa

> Observed MW: 50-54 kDa

Purification Method:

Protein A purification

CloneNo.: 1H10B3

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:20-1:200

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA Species Specificity: human, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

WB: IMR-32 cells, human colon tissue, rat brain tissue, rat colon tissue, mouse colon tissue, rat spinal cord

IHC: human brain tissue, human colon tissue

Background Information


Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3 $\,$

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66317-1-1g (Peripherin antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 66317-1-Ig (Peripherin antibody) at dilution of 1:100 (under 10x lens).




Immunohistochemical analysis of paraffinembedded human brain tissue slide using 66317-1-Ig (Peripherin antibody) at dilution of 1:100 (under 40x lens).



Immunohistochemical analysis of paraffinembedded human colon tissue slide using 66317-1- \lg (Peripherin Antibody) at dilution of 1:1000 (under 40x lens).



1X10^6 PC-12 cells were intracellularly stained with 0.4 ug Anti-Human Peripherin (66317-1-1g, Clone:1H10B3) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



rat brain tissue were subjected to SDS PAGE followed by western blot with 66317-1-1g (Peripherin Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.