For Research Use Only

SFRP2 Monoclonal antibody

Catalog Number:66328-1-lg 1 Publications



Basic Information

Catalog Number: 66328-1-lg

Isotype:

Size: 1000 µg/ml Source: Mouse

lgG1 Immunogen Catalog Number:

AG18840

295 aa, 34 kDa Observed MW:

38 kDa

BC008666

6423

GeneID (NCBI):

UNIPROT ID:

Q96HF1

Full Name:

Calculated MW:

GenBank Accession Number:

secreted frizzled-related protein 2

Positive Controls:

WB: pig brain tissue, Y79 cells, COLO 320 cells, rat brain tissue, rabbit brain tissue, mouse brain tissue

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:1000-1:6000 IHC 1:50-1:500

IF/ICC 1:750-1:3000

CloneNo.:

1F3A1

IHC: human malignant melanoma tissue,

IF/ICC: A549 cells,

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications:

WB

Species Specificity:

human, rat, pig, rabbit, mouse

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Secreted frizzled-related protein 2 (Sfrp2) is a secreted glycoprotein molecule containing an N terminal cysteinerich domain, which is typically 30-50% identical to the putative Wnt-binding site of the frizzled receptor, and a C terminal heparin-binding domain with weak homology to netrins. It has been implicated in diverse cellular processes, including embryogenesis, regulation of cell apoptosis, and cell differentiation. Methylation of this gene is a potential marker for the presence of colorectal cancer.

Notable Publications

Author **Pubmed ID** Journal Application Xin Tao 30954576 Exp Cell Res WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66328-1-lg (SFRP2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control




Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 66328-1-Ig (SFRP2 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human malignant melanoma tissue slide using 66328-1-Ig (SFRP2 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using SFRP2 antibody (66328-1-lg, Clone: 1F3A1) at dilution of 1:1500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



rat brain tissue were subjected to SDS PAGE followed by western blot with 66328-1-1g (SFRP2 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.