For Research Use Only

SMYD3 Monoclonal antibody

Catalog Number: 66330-1-Ig



Basic Information

Catalog Number: GenBank Accession Number: 66330-1-lg BC031010

 Size:
 GeneID (NCBI):
 CloneNo.:

 1700 μ g/ml
 64754
 185C10

 Source:
 HNURPOT ID:
 Recommon

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Mouse
 Q9H784
 WB 1:500-1:2000

 Isotype:
 Full Name:
 IHC 1:50-1:500

 IgG1
 SET and MYND domain containing 3
 IF/ICC 1:200-1:800

Immunogen Catalog Number: Calculated MW:

AG2624 428aa,49 kDa; 369aa,42 kDa

Observed MW: 49 kDa

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: A431 cells, COLO 320 cells, HEK-293 cells, MCF-7

Purification Method:

Protein G purification

cells

IHC: human pancreas cancer tissue, human liver

cancer tissue

IF/ICC : MCF-7 cells.

Background Information

SMYD3, also name as ZMYND1 and ZNFN3A1, belongs to the histone-lysine methyltransferase family. It is a histone methyltransferase that plays an important role in transcriptional regulation in human carcinogenesis. It can specifically methylate histone H3 at lysine 4 and activate the transcription of a set of downstream genes, including several oncogenes (e.g., N-myc, CrkL, Wnt10b, RIZ and hTERT) and genes involved in the control of cell cycle. (PMID: 20957523). It plays an important role in transcriptional activation as a member of an RNA polymerase complex. SMYD3 is frequently overexpressed in different types of cancer cells. It functions as a coactivator of Era and potentiates Era activity in response to ligand. SMYD3 as a new coactivator for ER-mediated transcription, providing a possible link between SMYD3 overexpression and breast cancer. (PMID: 19509295) The common variable number of tandem repeats polymorphism in SMYD3 is a susceptibility factor for some types of human cancer (PMID:16155568). Mainly cytoplasmic when cells are arrested at GO/G1, accumulates in the nucleus at S phase and G2/M(Uniprot).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed MCF-7 cells using SMYD3 antibody (66330-1-Ig, Clone: 1B5C10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), (CL594-Phalloidin, red).



A431 cells were subjected to SDS PAGE followed by western blot with 66330-1-1g (SMYD3 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 66330-1-lg (SMYD3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).