

For Research Use Only

IGSF2/CD101 Monoclonal antibody

Catalog Number: 66333-1-Ig



Basic Information

Catalog Number: 66333-1-Ig	GenBank Accession Number: BC136484	Purification Method: Protein A purification
Size: 1300 ug/ml	GeneID (NCBI): 9398	CloneNo.: 1E6A7
Source: Mouse	UNIPROT ID: Q93033	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:50-1:500
Isotype: IgG1	Full Name: immunoglobulin superfamily, member 2	
Immunogen Catalog Number: AG23297	Observed MW: 120 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls:
Species Specificity: human, mouse	WB : RAW 264.7 cells, NR8383 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human lung cancer tissue,

Background Information

IGSF2, also known as CD101, is a low abundance protein. Expression of CD101 is found on monocytes, granulocytes and dendritic cells. In addition expression on T lymphocytes is important for cell activation. It plays a role as inhibitor of T-cells proliferation induced by CD3 and inhibits expression of IL2RA on activated T-cells and secretion of IL2.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66333-1-Ig (IGSF2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66333-1-Ig (IGSF2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66333-1-Ig (IGSF2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



NR8383 cells were subjected to SDS PAGE followed by western blot with 66333-1-Ig (IGSF2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.