For Research Use Only

CLIC4 Monoclonal antibody

Catalog Number:66343-1-lg Featured Product

1 Publications



Basic Information

Catalog Number: 66343-1-lg

Size: 2000 ug/ml

Source:

Mouse

Isotype:

GenBank Accession Number: BC012444 GeneID (NCBI): 25932 **UNIPROT ID:** Q9Y696 Full Name:

lgG1 chloride intracellular channel 4 Calculated MW: Immunogen Catalog Number:

AG2942 253 aa, 29 kDa Observed MW:

29 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB. IF

Species Specificity: human, rat, pig **Cited Species:**

rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HuH-7 cells, pig heart tissue, HeLa cells, U2Os cells, pig kidney tissue, rabbit kidney tissue, rat kidney tissue, mouse kidney tissue, rat brain tissue, mouse brain tissue

Purification Method:

Protein G purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:50-1:500

IF/ICC 1:1000-1:4000

CloneNo.:

1B7F9

IHC: human heart tissue, human kidney tissue

IF/ICC: HeLa cells,

Background Information

Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transpithelial transport, maintenance of intracellular pH, and regulation $of cell\ volume.\ CLIC4 is\ a\ p53- and\ tumor\ necrosis\ factor\ alpha-regulated\ cytoplasmic\ and\ mitochondrial\ protein$ that belongs to the CLIC family of intracellular chloride channels. CLICs have actions distinct from traditional cell $membrane\ chloride\ channels,\ including\ formation\ of\ ion\ channels\ in\ intracellular\ organelles\ and\ roles\ in\ membrane\ order and\ o$ trafficking, apoptosis, and cell differentiation.

Notable Publications

Author	Pubmed ID	Journal	Application
Songhan Li	35498125	Oxid Med Cell Longev	WB,IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




WB result of CLIC4 antibody (66343-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CLIC4 transfected HeLa cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66343-1-lg (CLIC4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human heart tissue slide using 66343-1-Ig (CLIC4 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human heart tissue slide using 66343-1-Ig (CUC4 Antibody) at dilution of 1:200 (under 40x lens).



1x10^6 BT-549 cells were intracellularly stained with 0.2 μ g CLIC4 Monoclonal antibody (66343-1-1g, Clone: 1B7F9) and CoraLite® Plus 647-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO.RGAM005). Mouse 1gG1 isotype control (66360-1-1g, Clone: 1F8D3, blue) was parallel stained as control. Cells were fixed with 4% PFA.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using CLIC4 antibody (66343-1-lg, Clone: 1B7F9) at dilution of 1:2000 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).