For Research Use Only

COX2/ Cyclooxygenase 2/ PTGS2 Monoclonal antibody, PBS Only

Size: 1mg/ml



Purification Method:

Protein A purification

CloneNo.:

3G2B9

Catalog Number:66351-1-PBS Featured Product

Basic Information

Catalog Number:

66351-1-PBS

BC013734 GeneID (NCBI):

Source: **UNIPROT ID:** Mouse P35354 Isotype: Full Name: lgG2a prostaglandin-endoperoxide synthase

Immunogen Catalog Number:

AG24721

2 (prostaglandin G/H synthase and

GenBank Accession Number:

cyclooxygenase) Calculated MW: 604 aa, 68 kDa

Observed MW: 66-70 kDa

Applications

Tested Applications: WB, IHC, IF-P, ELISA Species Specificity: human, mouse

Background Information

COX2 (Prostaglandin G/H synthase 2, PTGS2) mediates the formation of prostaglandins from arachidonate. Its subunit structure is homodimer. The fully N-glycosylated PTGS2 is 72-74 kDa and the aglycosylated is 66 kDa (PMID:19656660). It also expresses a band of 39 kDa after unspecific cleavage (PMID:17509125). The 50 kDa band of fragmented PTGS2 has also previously been detected in AD brains (PMID:14724276).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data




Non-treated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 66351-1-Ig (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 66351-1-Ig (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66351-1-lg (COX2/ Cyclooxygenase 2/ PTGS2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse lung tissue using COX2/Cyclooxygenase 2/ PTGS2 antibody (66351-1-lg, Clone: 3G2B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66351-1-PBS in a different storage buffer formulation.