For Research Use Only

Cyclin D3 Monoclonal antibody, PBS Only



Catalog Number:66357-1-PBS


Featured Product

Basic Information	Catalog Number: 66357-1-PBS	GenBank Accession Number: BC011616	Purification Method: Protein A purification
	Size: 1 mg/ml	GenelD (NCBI): 896	CloneNo.: 1F2C6
	Source: Mouse	ENSEMBL Gene ID: ENSG00000112576	
	lsotype: lgG1	UNIPROT ID: P30281	
	Immunogen Catalog Number: AG22825	Full Name: cyclin D3	
		Calculated MW: 33 kDa	
		Observed MW: 33-35 kDa	
Applications	Tested Applications: WB,IHC,ELISA		
	Species Specificity: human		
Background Information	Cyclin D3 belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. The CDK4 activity associated with this cyclin was reported to be necessary for cell cycle progression through G2 phase into mitosis after UV radiation.		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS only		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 66357-1-1g (Cyclin D3 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66357-1-PBS in a different storage buffer formulation.



WB result of Cyclin D3 antibody (66357-1-Ig; 1:1500; incubated at room temperature for 1.5 hours) with sh-Control and sh-Cyclin D3 transfected Jurkat cells. This data was developed using the same antibody clone with 66357-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 66357-1-1g (Cyclin D3 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66357-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 66357-1-1g (Cyclin D3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66357-1-PBS in a different storage buffer formulation.