

For Research Use Only

NONO Monoclonal antibody

Catalog Number: 66361-1-Ig **1 Publications**



Basic Information

Catalog Number:

66361-1-Ig

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG17609

GenBank Accession Number:

BC010049

GeneID (NCBI):

4841

UNIPROT ID:

Q15233

Full Name:

non-POU domain containing,
octamer-binding

Calculated MW:

54 kDa

Observed MW:

54 kDa

Purification Method:

Protein G purification

CloneNo.:

2A2B10

Recommended Dilutions:

WB 1:500-1:2000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, rat, mouse

Cited Species:

mouse

Positive Controls:

WB: C2C12 cell, mouse skeletal muscle tissue,
ROS1728 cells

Background Information

Non-pou domain-containing octamer-binding protein (NONO), other named p54nrb, belongs to DBHS family, whose protein are predominately nuclear and involves in transcriptional and posttranscriptional gene regulatory functions as well as DNA repair. This family has roles in a wide range of biological processes, such as the regulation of circadian rhythm, carcinogenesis and progression of cancer. NONO forms heterodimer with SFPQ, and has been implicated in nuclear retention of A- to I-edited RNA as paraspeckle components, pre-mRAN 3'-end formation, cAMP cycling and transcriptional activation.

Notable Publications

Author	Pubmed ID	Journal	Application
Nian Liu	35953472	Nat Commun	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



C2C12 cell were subjected to SDS PAGE followed by western blot with 66361-1-Ig (NONO Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.