For Research Use Only

Occludin Monoclonal antibody

Catalog Number:66378-1-Ig Featured Product

133 Publications




Basic Information	Catalog Number: GenBank Accession Number: 66378-1-Ig BC029886		sion Number:	Purification Method: Protein G purification	
	Concentration:	GeneID (NCBI):		CloneNo.:	
	1000 ug/ml	4950		1D3C4	
	Source:	UNIPROT ID:		Recommended Dilutions:	
	Mouse	Q16625		WB: 1:5000-1:50000	
	lsotype: lgG1	Full Name: occludin		IHC: 1:50-1:500	
	Immunogen Catalog Number:	Calculated MW:			
	AG4057	522 aa, 59 kDa			
		Observed MW: 59 kDa			
Applications	Tested Applications:	sted Applications: Positive Controls:			
	WB, IHC, ELISA WB : HUVE		WB: HUVEC	cells, Jurkat cells, mouse colon tissue,	
			ls, U2OS cells, pig colon tissue, rat colon		
	WB, IHC tissue, 4T1 cells, HEK-293 cells				
			colon cancer tissue, mouse kidney tissu		
	human, mouse, rat, pig Cited Species:				
	human, mouse, rat, pig, bovine, sheep				
	Note-IHC: suggested a with TE buffer pH 9.0; (antigen retrieval may b with citrate buffer pH 6	*) Alternatively be performed	, ,		
Background Information	formation and regulation of the t	ellular N and C termir tight junction paracell s at the posttranscript	ni and two extracellu lular permeability b ional and posttransl	s a tetraspanin protein with four Ilar loops. Occludin plays a role in the arrier. Occludin can exist in different ational levels, the monomeric occludin	
Notable Publications	Author	Pubmed ID	Journal	Application	
	Dayun Feng	36179025	Sci Adv	WB	
	Yingying Wang	36174863	Int J Biol Macromol	WB	
	Cantao Yang	36108982	Food Chem Toxicol	WB	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66378-1-lg (Occludin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of Occludin antibody (66378-1-Ig; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-Occludin transfected Jurkat cells.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66378-1-1g (Occludin antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66378-1-1g (Occludin antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).