

For Research Use Only

CD80 Monoclonal antibody

Catalog Number: 66406-1-Ig **67 Publications**



Basic Information

Catalog Number: 66406-1-Ig	GenBank Accession Number: BC042665	Purification Method: Protein G purification
Concentration: 1500 ug/ml	GeneID (NCBI): 941	CloneNo.: 1E2F10
Source: Mouse	UNIPROT ID: P33681	Recommended Dilutions: WB 1:1000-1:4000
Isotype: IgG1	Full Name: CD80 molecule	
Immunogen Catalog Number: AG5615	Calculated MW: 33 kDa	
	Observed MW: 68 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Raji cells, DC2.4 cells, Daudi cells
Cited Applications: WB, IF	
Species Specificity: human, mouse, pig	
Cited Species: human, mouse, rat, pig, rabbit	

Background Information

CD80 (also known as B7-1) is a type I membrane protein that is a member of the immunoglobulin superfamily, with an extracellular immunoglobulin constant-like domain and a variable-like domain required for receptor binding. It is expressed on antigen-presenting cells (APCs), including B cells, dendritic cells, monocytes, and macrophages. CD80 is the receptor for the proteins CD28 and CTLA-4 found on the surface of T-cells. It is involved in the costimulatory signal essential for T-lymphocyte activation. T-cell proliferation and cytokine production is induced by the binding of CD28, binding to CTLA-4 has opposite effects and inhibits T-cell activation. CD80 also acts as a cellular attachment receptor for adenovirus subgroup B. (PMID: 7545666; 12015893; 16920215)

Notable Publications

Author	Pubmed ID	Journal	Application
Yan Yang	34582523	Biomater Sci	
Chenxia Juan	30243953	Int J Biochem Cell Biol	WB
Yuanhang Li	30411299	Pathol Oncol Res	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Raji cells were subjected to SDS PAGE followed by western blot with 66406-1-Ig (CD80 / B7-1 antibody at dilution of 1:2000 incubated at room temperature for 1.5 hours.