### For Research Use Only

# ZC3HAV1 Monoclonal antibody

Catalog Number:66413-1-lg Featured Product

4 Publications



**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:1000-1:6000 IHC 1:50-1:500

IF/ICC 1:200-1:800

CloneNo.:

1G10B9

**Basic Information** 

Catalog Number: GenBank Accession Number: 66413-1-lg BC025308 GeneID (NCBI): Size: 1400 ug/ml 56829 **UNIPROT ID:** Source: Mouse Q7Z2W4 Full Name: Isotype:

lgG1 zinc finger CCCH-type, antiviral 1 Calculated MW: Immunogen Catalog Number:

AG10364 699aa,78 kDa; 902aa,101 kDa

> Observed MW: 100 kDa

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications:

WB, IF

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, HEK-293 cells, HeLa cells

IHC: human kidney tissue, IF/ICC: HepG2 cells,

# **Background Information**

he zinc-finger antiviral protein (ZAP) was originally identifed from a rat cDNA library due to its conferring resistance to retroviruses. ZC3HAV1 is a CCCH-type zinc finger protein, it may primarily function to inhibit viral gene  $expression\ and\ induce\ an\ innate\ immunity\ to\ viral\ infection.\ Alternative\ splicing\ occurs\ at\ this\ locus\ and\ two$ variants, each encoding distinct isoforms.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ana Cristina Gonzalez-Perez	33947766	mBio	WB
Marlene V Buckmaster	35913217	J Virol	IF
Zehua Shang	38750080	Nat Commun	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



250 kDa→
150 kDa→
100 kDa→
70 kDa→

WB result of ZC3HAV1 antibody (66413-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZC3HAV1 transfected HEK-293 cells.

Jurkat cells were subjected to SDS PAGE followed by western blot with 66413-1-lg (ZC3HAV1 Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66413-1-Ig (ZC3HAV1 Antibody) at dilution of 1:300 (under 40x lens).

Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using ZC3HAV1 antibody (66413-1-lg, Clone: 1G10B9) at dilution of 1:400 and MultirAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).