For Research Use Only

## LDLR Monoclonal antibody

Catalog Number:66414-1-lg Featured Product 7 Publications







Basic Information	Catalog Number: 66414-1-lg	GenBank Accession Numb BC014514	per: Purification Method: Protein G purification	
	Concentration:	GenelD (NCBI):	CloneNo.:	
	1200 ug/ml	3949	1A2A3	
	Source: Mouse	ENSEMBL Gene ID: ENSG00000130164	Recommended Dilutions: WB 1:5000-1:15000	
	Isotype:	UNIPROT ID:	IHC 1:50-1:500	
	lgG1	P01130		
	Immunogen Catalog Number: AG1236	Full Name: low density lipoprotein receptor		
		Calculated MW: 95 kDa		
		Observed MW: 120 kDa, 160 kDa		
Applications	Tested Applications:	Positive Controls:		
	Cited Applications:cells, MCF-7 cells, Raji ceWBIHC : human liver tissue,		B : HeLa cells, HEK-293 cells, Jurkat cells, HepG Ils, MCF-7 cells, Raji cells	
			C : human liver tissue,	
	Species Specificity: human			
	Cited Species: human, hamster			
	Note-IHC: suggested antige TE buffer pH 9.0; (*) Alterr retrieval may be performe buffer pH 6.0	atively, antigen		
Background Information	LDLR (low density lipoprotein receptor) is a member of the LDL receptor gene family and is involved in receptor- mediated endocytosis of specific ligands. The LDLR is a cell surface glycoprotein that scavenges LDL from the blo and regulates plasma LDL cholesterol. The cytoplasmic domain of the LDL receptor is necessary for the receptor to cluster in coated pits, which promotes the rapid endocytosis of bound LDL. The protein is highly glycosylated thro N- and O-linkages and thus migrates at 100 to 160 kDa bands on SDS-PAGE.			
Notable Publications	Author	Pubmed ID Journal	Application	
	Zoe White	36447272 Skelet Mu		
	Jun Xiao	30864739 Int J Mol N	Aed WB	
	Shui-Mei Sun	32724176 Acta Phar	macol Sin WB	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide an Aliquoting is unnecessary for -20	d 50% glycerol, pH7.3		

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




WB result of LDLR antibody (66414-1-lg; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-LDLR transfected HeLa cells.



Various lysates were subjected to SDS PAGE followed by western blot with 66414-1-lg (LDLR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66414-1-Ig (LDLR Antibody) at dilution of 1:400 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66414-1-Ig (LDLR Antibody) at dilution of 1:400 (under 40x lens).