

For Research Use Only

Amphiregulin Monoclonal antibody

Catalog Number: 66433-1-Ig **3 Publications**



Basic Information

Catalog Number: 66433-1-Ig	GenBank Accession Number: BC009799	Purification Method: Protein A purification
Size: 1500 µg/ml	GeneID (NCBI): 374	CloneNo.: 1A1G9
Source: Mouse	UNIPROT ID: P15514	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:50-1:500
Isotype: IgG1	Full Name: amphiregulin	
Immunogen Catalog Number: AG8907	Calculated MW: 252 aa, 28 kDa	
	Observed MW: 50 kDa, 37 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : A549 cells, rat brain tissue, MCF-7 cells, pig brain tissue IHC : human pancreas cancer tissue, human colon cancer tissue
Cited Applications: IF, WB	
Species Specificity: human, rat, pig	
Cited Species: human	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Amphiregulin (AREG) is one of the ligands of the epidermal growth factor receptor (EGFR). AREG plays a central role in mammary gland development and branching morphogenesis in organs and is expressed both in physiological and in cancerous tissues. The AREG protein is synthesized as a 252-amino acid transmembrane precursor, pro-AREG. At the plasma membrane, pro-AREG is subjected to sequential proteolytic cleavages within its ectodomain and is then released as the soluble AREG protein. Depending on the cell type and microenvironment, AREG can be produced in multiple cellular and mature forms using alternative pro-AREG cleavage sites and glycosylation motifs. Post-translational modifications of 50-kDa pro-AREG produces a major soluble 43-kDa form, 28-, 26-, 16-kDa membrane anchored forms, and soluble 21-, 19-, and 9-kDa forms (PMID: 9642297).

Notable Publications

Author	Pubmed ID	Journal	Application
Jie Liu	30745837	Int J Biol Sci	
Yingjian Huang	34358528	J Invest Dermatol	WB
Yiyun Chen	37652019	Cell Rep Med	IF


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A549 cells were subjected to SDS PAGE followed by western blot with 66433-1-Ig (Amphiregulin antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66433-1-Ig (Amphiregulin antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66433-1-Ig (Amphiregulin antibody) at dilution of 1:300 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human pancreas cancer tissue slide using 66433-1-Ig (Amphiregulin antibody) at dilution of 1:300 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).